Effect of arbuscular mycorrhiza fungi and organic fertilizers on yield and nutrients uptake of two wheat cultivars

B. Kermanizadeh¹, A. Gholamalizadeh Ahangar^{1*}, S. K. Sabbagh² and A. Sirousmehr³

(Received: 10 June 2013; Accepted: 3 June 2015)

Abstract

This research was conducted in order to evaluate the direct effects of organic and biofertilizers on yield of two native wheat cultivars (Bolani and cross-Bolani) in Sistan area. The experiment was performed as a factorial, based on a completely randomized design with three replications, in Research Greenhouse of University of Zabol. In this study, fertilizer factor at 8 levels [Vermicompost (F1), vermicompost + compost (F2), vermicompost + mycorrhiza (F3), vermicompost + mycorrhiza + compost (F4), compost (F5), compost + mycorrhiza (F6), mycorrhiza (F7) and control (no fertilizer application, F8)] and two wheat cultivars [Bolani (C1) and cross-Bolani (C2)] were considered. Results showed that the highest grain yield (1.13 g/pot) was obtained from combination of mycorrhiza and cross-Bolani treatments (F7C2). Combined treatments of compost + mycorrhiza and cross-Bolani (F6C2) and vermicompost + compost and cross-Bolani (F2C2) were more suitable for nitrogen uptake. The highest percentage of protein (10.27%) was resulted from F6C2 and F2C2 treatment. Overall, combined treatments of F6C2 and F2C2 seem appropriate for Bolani and cross-Bolani wheat cultivars.

Keywords: Wheat, Organic fertilizer, Vermicompost, Mycorrhiza, Compost.

^{1.} Dept. of Soil Sci., College of Agric., Univ. of Zabol, Zabol, Iran.

^{2.} Dept. of Plant Protec., College of Agric., Univ. of Zabol, Zabol, Iran.

^{3.} Dept. of Agron. and Plant Breed., College of Agric., Univ. of Zabol, Zabol, Iran.

^{*:} Corresponding Author, Email: ahangar@uoz.ac.ir