

## Comparison of yield, yield components and seed quality (oil and protein content) of two rapeseed cultivars as affected by different levels of soil-applied nitrogen and zinc

V. Olama<sup>1\*</sup>, A. M. Ronaghi<sup>1</sup>, N. A. Karimian<sup>1</sup>, J. Yasrebi<sup>1</sup>,  
R. Hamidi<sup>2</sup> and M. Tavajjoh<sup>1</sup>

(Received: 23 Sep-2012 ; Accepted: 15 Apr-2013)

### Abstract

A greenhouse experiment was conducted in order to evaluate the influence of different nitrogen (N) and zinc (Zn) levels on yield, yield components, seed quality and N and Zn contents of two cultivars of rapeseed (*Brassica napus* L.). In this study, two cultivars of rapeseed (Talayeh and Opera), three N levels (100, 200 and 300 mg/kg) and three Zn levels (0, 5 and 10 mg/kg) with three replications were used. The results showed that significant differences were observed between the cultivars in the studied traits. Increasing N and Zn significantly increased plant height, number of siliques per plant, number of seeds per silique, 1000-seed weights, seed yield, oil, straw and root yields and protein content. Oil seed content was the highest in application of 100 mg/kg N rate and then decreased or increased at higher N and Zn rates, respectively. The highest grain yield, oil yield and protein percentage were obtained with 300 mg/kg N and 10 mg/kg Zn. Increase in seed yield was related to increase in the number of siliques per plant and number of seeds per silique. Application of treatments increased concentration of N and Zn in the seeds, straw, and roots of both rapeseed cultivars. The interaction of N and Zn showed significant effects on most studied traits of rapeseed.

**Keywords:** Oil seeds, Qualitative traits, Protein Oil.

---

1. Dept. of Soil Sci., College of Agric., Shiraz Univ., Shiraz, Iran.

2. Dept. of Agron. and Plant Breed., College of Agric., Shiraz Univ., Shiraz, Iran.

\*: Corresponding Author, Email: vidaolama@gmail.com