Effect of organic and inorganic substrates on vegetative and ecophysiological characteristics of coleus (*Coleus blumei*)

A. Sajjadinia¹*, M. Khayyat² and H. R. Karimi³

(Received : 5 December 2010 ; Accepted : 6 March 2011)

Abstract

In order to study of the effect of substrates on vegetative and ecophysiological characteristics of coleus plant, coleus cuttings containing two leaves were planted in perlite, cocopeat, peatmoss, compost, 50% perlite + 50% cocopeat, 50% perlite + 50% cocopeat, 50% perlite + 50% cocopeat, 50% perlite + 50% compost and 50% peatmoss, 50% compost with four replicates. Cuttings were irrigated by hydroponic solution after root formation. Results indicated that the highest root fresh and dry weight was observed in 50% perlite + 50% compost treatment, the highest secondary roots in perlite treatment, and the longest root and stem, the most fresh and dry weight of shoot, number of leaves and photosynthesis were observed in peatmoss treatment. Result showed that the best substrate for growth and development of coleus plant is peatmoss.

Keywords: Peatmoss, Perlite, Coleus, Cocopeat, Substrate.

^{1.} MSc., Dept. of Hort., Faculty of Agric., Vali-e-Asr Univ. of Rafsanjan, Iran.

^{2.} Dept. of Hort., Faculty of Agric., Birjand Univ., Birjand, Iran.

^{3.} Assist. Prof., Dept. of Hort., Faculty of Agric., Vali-e-Asr Univ. of Rafsanjan, Iran.

^{*:} Corresponding Author, Email: sajjadinia@yahoo.com